

Fludinal®

Your doctor prescribed Fludinal®	oral solution with a	formula that contains:
Each 5 mL contains:		
Ambroxol HCI		15 mg
Vehicle gs.		

What are the indications for Fludinal®? In is indicated for patients with acute bronchitis, chronic bronchitis, acute exacerbation of bronchitis with presence of bronchial hyperactivity, bronchial asthma, cystic fibrosis, bronchiectasis, sinusitis and all those pathologies with bronchial alteration.

What is the dose recommended for Fludinal®? The best dose is the one indicated by your doctor but as reference dose we recommend for children aged 2 to 5 years: one half teaspoon (2.5 mL) every 8 hours; children aged 5 to 12 years: 1 teaspoon (5 mL) cor 3 time a day; adults and children older than 12 years of age: 2 teaspoons (10 mL) 3 times a day for the first two days, and then 1 teaspoon (5 mL) 2 or 3 times a day.

What are the contraindications of Fludinal®? Hypersensitivity to any of the formula's components. Children under 2 years of age.

What precautions must I take when taking Fludinal®? Pregnancy and breastfeeding. It contains phenylalanine, caution is advised in patients with phenylketonuria.

There have been reports of serious skin reactions such as erythema multiforme, Stevens-Johnson syndrome (SSJ), Toxic Epidermal Necrolysis (NET), Acute Generalized Exanthematous Pustulosis (PEGA) associated with the administration of Ambroxol. If the patient has symptoms or sings of a progressive rash (sometimes associated with blisters or mucosal lesions), treatment with ambroxol should be stopped immediately and the doctor should be consulted.

What is the presentation of Fludinal®? 120 mL bottle in a box.

FOLLOW YOUR DOCTOR'S DIRECTIONS EXACTLY, CONSULTING HIM/HER PERIODICALLY AND IN RETURN YOU WILL GET GOOD RESULTS

Pharmalat, S. A.
0 Avenida "C" 2-55, Colonia Najarito
Zona 6 de Villa Nueva, Guatemala, C. A.
PBX: (502) 6628-1100 • FAX: (502) 6628-1157

 $www.pharmalat.net\\servicioal cliente@pharmalat.com.gt$



