



Molly

Against fever and pain **ACETAMINOPHEN DROPS** Strawberry flavor

Drug facts

Active ingredients:

Drops: Each mL contains: Paracetamol (Acetaminophen).....100 mg

Indications and uses: Analgesic and antipyretic effective against mild to moderate pain caused by joint conditions, earache, headache, toothache, neuralgia, minor surgical procedures, etc. It is also effective for the treatment of fever caused by viral infections, fever post vaccination, etc.

Dose:

- Children from 0 to 3 months: 0.6 mL equivalent to 15 drops every 6 hours
- Children from 4 to 11 months: 1.2 mL equivalent to 30 drops every 6 hours
- Children from 12 to 23 months: 1.6 mL every 6 hours
- Children from 2 to 3 years old: 2 mL every 6 hours
- Children from 4 to 5 years old: 2.8 mL every 6 hours

Contraindications:

- Hypersensitivity to any component of the formula.
- Administer with caution in patients with liver damage or bleeding from the gastrointestinal tract, should not be administered for prolonged periods.
- In pregnant or lactating should consult your doctor before use.

General precautions:

- If a rare sensitivity reaction occurs, discontinue the drug immediately .
- If fever or pain persists seek medical advice.
- Do not use higher doses than indicated.

Warnings: Do not administer the recommended dose more than five times a day, or for more than five consecutive days to children, or for more than ten consecutive days to adults. If pain or fever persists for more than three consecutive days, consult a doctor.

Interactions: The potential hepatotoxicity of Paracetamol can be increased by the administration of large doses or prolonged doses of a series of substances, by induction of hepatic microsomal enzymes. These agents can also cause a decrease in the therapeutic effects of Paracetamol. These substances are: Barbiturics, Carbamazepine, Isoniacide, Hydantoins, Rifampicin and Sulfinpyrazone.

Adverse reactions: Adverse reactions of Paracetamol are usually rare or very rare; among which may be, malaise, increased levels of liver transaminases and hypotension.

Overdose: Symptoms of overdose include dizziness, vomiting, loss of appetite, jaundice, abdominal pain, and kidney and liver failure. If an overdose has been ingested, the patient should be treated promptly in a medical center, even if there are no significant symptoms or signs, since, although these can cause death, they often do not manifest immediately after ingestion, but rather after third day. Death due to hepatic necrosis may occur. Likewise, acute renal failure can appear.

Route of administration: oral.

Presentation: Dropper bottle 10, 15, 20, 25 and 30 mL.

Storage conditions:

- Keep in a cool and dry place (below 30°C), protect from light.
- Medicine, keep out of reach of children.