



Iza

Eliminate parasites

ALBENDAZOLE

Oral suspension

400 mg / 10 mL | Orange flavor

Drug facts

Active Ingredient: each 10 mL contains:

Albendazole.....400 mg
Vehicle q.s.

Indications and uses:

Albendazole is effective for treating infections caused by *Ascaris lumbricoides*, *Trichuris trichiura*, *Enterobius vermicularis*, *Ancylostoma duodenale*, *Necator americanus*, *Hymenolepis nana*, *Taenia* sp, *Strongyloides stercoralis*, *O. viverrini*, *C. sinensis*, cutaneous larva migrans and *G. spinigerum*, also active against *Giardia lamblia*. It is also indicated for the treatment of parenchymal neurocysticercosis due to active lesions caused by larval forms of the pork tapeworm, *Taenia solium*. It is also useful for treatment of hydatid disease of the liver, lungs and peritoneum caused by the larval form of the dog tapeworm, *Echinococcus granulosus*.

Dose:

- Children over 2 years and adults: 10 mL suspension (400 mg) in one shot.
- In estrongliodiasis, *Taenia*, or infections by *Himenolepis*, 10 mL suspension in a single dose for 3 consecutive days. No fast or use of laxatives is necessary. Always consult your doctor.

Shake well before use.

Contraindications:

- Hypersensitivity to any component of the formula.
- Is contraindicated in patients with known benzimidazole-type compounds, or to any component of the formulation Albendazole hypersensitivity.

Warnings:

- Consult your doctor before starting to use this medicine if you are pregnant or suspect that you are pregnant or while breast-feeding.

General precautions:

- If a rare sensitivity reaction occurs, discontinue the drug immediately.
- Patients who are being treated for neurocysticercosis should receive adequate anticonvulsant and steroid therapy as required. Consideration should be given to oral or intravenous corticosteroids to prevent cerebral hypertensive episodes during the first week of treatment anticysticercosis.
- Cysticercosis may rarely involve the retina. If injuries are displayed, therapy against cysticercosis should be evaluated against the potential damage to the retina, due to the administration of Albendazole.

Administration way: oral.

Presentation:

10 mL bottle.

Storage conditions:

- Store in a cool, dry place at room temperature not exceeding 30°C, protected from light.
- Keep out of reach of children.