



AMIBEST® COATED TABLETS

COMPOSITION:

Each coated tablet contains:

Secnidazole 500 mg

Excipients qs

INDICATIONS:

Urethritis and vaginitis produced by *Trichomonas vaginalis*. Giardiasis, intestinal and hepatic amebiasis.

CONTRAINDICATIONS:

Hipersensitivity to secnidazole, blood dyscrasias, pregnancy and breast-feeding.

DOSAGE AMIBEST® coated tablets:

Urethritis and vaginitis produced by *Trichomonas vaginalis*: 4 tablets (2 g) as a single dose in the morning.

Intestinal amebiasis and giardiasis: 2 tablets (1 g) in the morning and 2 tablets (1 g) in the evening, in treatment of one day.

Hepatic amebiasis: 3 tablets (1.5 g) as a single dose or divided in several takes along the day, during 5 days.

PACKINGS:

Box with 8 coated tablets in blister.

Fabricado en Guatemala por
Pharmalat, S. A. para
Winzzer Medical Corporation
0 Avenida "C" 2-55, Colonia Najarito
Zona 6 de Villa Nueva, Guatemala, C. A.
PBX: (502) 6628-1100 • FAX: (502) 6628-1157
www.pharmalat.net
servicioalcliente@pharmalat.com.gt

