



GESIMAX[®]

NAPROXEN SODIUM

COMPOSITION:

Each tablet contains:

Naproxen sodium550 mg

Excipients qs

Each 5 mL oral suspension contains:

Naproxen sodium125 mg

Vehicle qs

INDICATIONS: Mild to moderate pain, acute or chronic conditions with or without inflammation, fevers, trauma and musculoskeletal disorders and of the joints. Reumateudea, arthritis, osteoarthritis, ankylosing, dysmenorrhea, acute gout.

CONTRAINDICATIONS: Hypersensitivity to naproxen or other non steroid antiinflammatories (NSAID's), patients with peptic ulcer, renal or hepatic failure, pregnancy and breast-feeding.

INTERACTIONS: May potentiate the action of alcohol an toxicity, methotrexate, digoxin, lithium salts, antidiabetics, oral anticoagulants and valproic acid, it reduces effect of diuretics beta blockers and ACE inhibitors.

SIDE EFFECTS: Edema, nausea, abdominal pain, constipation, flatulence, headache, dizziness, blurred vision, drowsiness, tinnitus, facial inflammation, asthma and shock.

PRECAUTIONS AND WARNINGS: In patients with gastrointestinal disease administer under close medical surveillance. The administration of this product may unblance the diet in patients who follow a low sodium regime.

DOSAGE TABLETS: Adolescents and adults: 1 tablet 2 times per day until desappearance of symptoms or according to the physians indication.

DOSAGE ORAL SUSPENSION: Children: 10 mg/Kg body weight, an initial dose followed by 2.5 to 5 mg/Kg body weight every 8 hours. The dose should not exceed 15 mg/Kg per day after ther first day of treatment.

PACKINGS:

Pack with 10 tablets in blister and dispenser with 100 tablets in blister.

Oral suspension bottle of 60mL (shake well before use).

0 Avenida "C" 2-55, Colonia Najarito
Zona 6 de Villa Nueva, Guatemala, C. A.
PBX: (502) 6628-1100 • FAX: (502) 6628-1157
www.pharmalat.net
servicioalcliente@pharmalat.com.gt

